

October 27, 2026 - Dr. Krebs (claudia.krebs@ubc.ca)




Objectives:

1. Identify the basal nuclei (basal ganglia), thalamus, internal capsule, and associated structures on brain specimens, coronal and horizontal sections, and relevant clinical images, and describe their anatomical relationships.
2. Trace the major connections of the direct and indirect basal ganglia pathways and predict how each pathway influences thalamocortical activity and movement. Relate the net effect of the direct and indirect pathways on the thalamus to their role in movement.
3. Relate motor, associative, and limbic basal ganglia circuits to their corresponding cortical regions and functions.
4. Identify the principal arterial supply to the basal nuclei and internal capsule, and use vascular territories and anatomical relationships to predict the effects of focal lesions.
5. Identify the major external and internal features of the cerebellum—including its lobes, deep nuclei, and peduncles—and relate the vestibulocerebellum, spinocerebellum, and cerebrocerebellum to their anatomical locations.
6. Trace the major afferent and efferent pathways through the cerebellar peduncles and explain how the functional divisions of the cerebellum contribute to balance, coordination, motor control, and motor learning. Describe the major tracts travelling in the cerebellar peduncles.
7. Identify the principal arterial supply to the cerebellum, and relate the vascular territories and anatomical relationships of the cerebellum to characteristic patterns of neurological dysfunction.
8. Integrate the clinical presentation, neurological findings, and imaging features with the underlying functional neuroanatomy to explain and localize a lesion to the basal ganglia or cerebellum.

Resources

Here are the e-tutorials, videos, and 3D specimens to prepare for this lab – click the green buttons to access them.



Modules:



Videos:



3D Models:



October 27, 2026 - Dr. Krebs (claudia.krebs@ubc.ca)

Overview of the Basal Ganglia

Glossary and Abbreviations:

- caudate nucleus + putamen = neostriatum or (dorsal) striatum
- putamen + globus pallidus = lentiform or lenticular nucleus
- globus pallidus, external segment (GPe)
- globus pallidus, internal segment (GPi)
- substantia nigra (SN)
- subthalamic nucleus (STN)

Micrographs

Coronal Sections

Horizontal Sections

On gross specimens & micrographs, identify:

- ☐ Putamen
- ☐ Globus pallidus (external & internal)
- ☐ Caudate nucleus
- ☐ Nucleus accumbens
- ☐ Substantia nigra
- ☐ Subthalamic nucleus (*micrographs only*)
- ☐ Ansa lenticularis & subthalamic fasciculus (*micrographs only*)
- ☐ Relationship of basal ganglia to internal capsule & thalamus


The inhibitory projection from the GPi to the thalamus is called ansa lenticularis.

Find this structure in your micrographs.



Which neurotransmitter is used by this projection?

Hide Labels

Coronal section of brain





October 27, 2026 - Dr. Krebs (claudia.krebs@ubc.ca)




Hide Labels

October 27, 2026 - Dr. Krebs (claudia.krebs@ubc.ca)



Hide Labels



C/P = Caudate and Putamen; GPe = external segment of Globus Pallidus; GPi = internal segment of Globus Pallidus; SN = Substantia Nigra; STN = Subthalamic Nucleus


- The **thalamus** is under tonic inhibition — GABAergic neurons from the GPi to the thalamus, traveling in the thalamic fasciculus.
- The **direct pathway** releases the thalamus from its tonic inhibition by inhibiting the GPi output. This inhibition comes through a direct projection from the striatum (C/P). This leads to increased output from the thalamus to the cortex and subsequently an increase in cortical output.
The direct pathway facilitates target oriented, efficient movement.
- The **indirect pathway** increases the inhibition of the thalamus by stimulating the GPi. The subthalamic nucleus (STN) stimulates the GPi, which in turn inhibits the thalamus. This leads to reduced output from the thalamus to the cortex and subsequently a decrease in cortical output.
The indirect pathway inhibits superfluous and potentially competing movements.
- Dopamine** facilitates cortical output and movement. It strengthens the direct pathway and weakens the indirect pathway. Dopaminergic projections come from the substantia nigra (SN).

October 27, 2026 - Dr. Krebs (claudia.krebs@ubc.ca)


Use these templates to draw in nuclei of the basal ganglia as seen in horizontal & coronal cross-sections.

Coronal Sections

Horizontal Sections



Horizontal





Coronal

October 27, 2026 - Dr. Krebs (claudia.krebs@ubc.ca)

Blood Supply to the Basal Ganglia

ACA (Deep Penetrating Branches)	• Caudate head • Putamen anterior • Globus pallidus anterior
MCA (Lenticulostriate Arteries)	• Caudate body • Putamen • Globus pallidus
PCA (Deep Penetrating Branches)	• Caudate (minor supply) • Substantia nigra • Subthalamic nucleus
AChA (from Internal Carotid)	• Caudate tail • Putamen posterior • Globus pallidus posterior





Krebs et al., Lippincott's Illustrated Reviews - Neuroscience, 2nd ed. Copyright © 2017 LWW. All rights reserved.

October 27, 2026 - Dr. Krebs (claudia.krebs@ubc.ca)

The Cerebellum

On gross specimens & micrographs, identify:

- ☐ Cerebellar hemispheres
- ☐ Cerebellar lobes
- ☐ Flocculus, nodule, vermis, cerebellar tonsils
- ☐ Dentate nucleus (micrographs only)
- ☐ Cerebellar peduncles




Hide Labels

October 27, 2026 - Dr. Krebs (claudia.krebs@ubc.ca)

Blood Supply to the Cerebellum

AICA	• Cerebellar peduncles • Flocculus, nodule (floconodular lobe) • Lateral aspect of posterior lobe on inferior surface • Assists with supply to deep cerebellar nuclei
PICA	• Posterior lobe: inferior surface, extending onto superior surface • Tonsils • Inferior part of vermis
SCA	• Anterior lobe • Posterior lobe: superior aspect, strip adjacent to anterior lobe • Vermis on superior surface • Major supply to deep cerebellar nuclei • Can assist with supply to superior cerebellar peduncle (SCP), rostral parts of middle cerebellar peduncle (MCP)




Krebs et al., Lippincott's Illustrated Reviews - Neuroscience, 2nd ed. Copyright © 2017 LWW. All rights reserved.

October 27, 2026 - Dr. Krebs (claudia.krebs@ubc.ca)


Blood Supply to the Cerebellum

On gross specimens & micrographs, identify:

- ☐ Posterior inferior cerebellar artery (PICA)
- ☐ Anterior inferior cerebellar artery (AICA)
- ☐ Superior cerebellar artery (SCA)



Krebs et al., *Lippincott's Illustrated Reviews - Neuroscience*, 2nd ed.
Copyright © 2017 LWW. All rights reserved.



Hide Labels

Vessels of posterior
brainstem and cerebellum

October 27, 2026 - Dr. Krebs (claudia.krebs@ubc.ca)

CASE 1

A **72-year-old man** presents to your office for a 6-month follow-up appointment.

You have been seeing him in regard to a **right hand tremor with slowness in movement on the right**.

He has had these symptoms for 9 years.

He is on medication for this concern.

He tells you that the tremor is getting worse and that it is more prominent when he is sitting at rest or walking.

His wife has noticed that his gait has been slow for the past 6 years.

He has also noticed that he is slower with movement in his right hand despite the medications that he is taking for this medical condition.

He denies any weakness or sensory changes.

He denies any unsteadiness except occasionally when he turns quickly he can lose his footing.

On review of his records, you recall the following trouble:

Smell: Yes, my food tastes bland, occurring for the past 10 years

Sleep: Yes, my wife tells me that I kick around in my sleep and appear distressed during this period

Urination: Yes, I am getting up 3-5 times in the middle of the night but no trouble during the day

Bowel function: Yes, I am constipated at times

Lightheadedness when standing: Occasionally, but only in the morning

Today you ask the same questions and find out that all the above positive symptoms have worsened in the past 6 months.

He is frequently lightheaded when standing and he is constipated all the time.

In addition you ask the following:

Behaviour changes: No

Mood concerns: He denies a low mood but his wife later tells you that she is concerned about his mood; he is less interactive, socially withdrawn, but there is no suicidal ideation.

Cognitive symptoms: He has noticed trouble with his ST memory of the past few months. He is not as quick to recall information and has word finding difficulties. He denies any geographic disorientation or impairment in judgement, insight or planning. He is able to do his ADL and IADLs. His friends have noticed this change.

On examination, his blood pressure lying down is 130/78 and standing is 104/72.

He has less animation in his face with reduced blinking.

He has rigidity on the right >> left arm with cogwheeling at both wrists.

There is a resting tremor in the right hand.


His strength, reflexes, sensation, and coordination is normal.

He has slow small amplitude movement of the right >>> left hand.

He has reduced arm swing on the right and slow gait with slight stooped posture.

He turns slowly.

October 27, 2026 - Dr. Krebs (claudia.krebs@ubc.ca)




What is this patient's medical condition?

Where is the pathology to explain why the right side is more involved?

What is the reason for the constipation and nocturia?

Why are his cognition and mood changing?




October 27, 2026 - Dr. Krebs (claudia.krebs@ubc.ca)

CASE 2

A **78-year-old man** with a history of **hypertension and diabetes** presents to your office for follow-up 2 months after his **stroke**.

You review the imaging and see that the stroke affected the **left subthalamic nucleus**.



Which vessel was likely involved in this stroke?

What type of movement disorder will this patient have? Why?

October 27, 2026 - Dr. Krebs (claudia.krebs@ubc.ca)



On which side will the movement disorder present? Explain why.